

DOCTORS FOSTER^{and}SMITH[®] PHARMACY

Patient Information Sheet

PO Box 100, Rhinelander WI 54501
For questions or to consult with a pharmacist, call 1-800-447-3021 Mon-Fri from 9am-6pm CST and Sat from 9am-1pm CST. If you have an emergency situation, please contact your local veterinarian immediately.

Desoxycorticosterone Pivalate (DOCP)

(Percorten-V)

Common Drug Name

Desoxycorticosterone pivalate (DOCP)

Common Brand Names

Percorten-V

No generic products are available.

Storage

Store at room temperature in a tight, light resistant, childproof container. Protect from freezing. Do not mix with any other injectable medication.

Uses

DOCP is a mineralocorticoid used to treat Addison's disease (adrenocortical insufficiency) in dogs and cats.

Dose and Administration

Always follow the dosage instructions provided by your veterinarian. If you have difficulty giving the medication, contact your veterinarian.

Dosage varies per patient and is adjusted based upon blood electrolyte (e.g., sodium and potassium) levels.

DOCP is an injectable medication. Shake the vial vigorously before each use, and follow your veterinarian's directions on how to give the injection. Injections are usually given every 21-30 days. Treatment is needed for life.

If you miss a dose, give it as soon as you remember. If it is almost time for the next dose, skip the one you missed and go back to the regular schedule. Do not give 2 doses at once.

This medication should only be given to the pet for whom it was prescribed.

Possible Side Effects

May see irritation at injection site. If given at too low of a dose, may see signs of Addison's disease (weakness, depression, lack of appetite, vomiting, and diarrhea). If given at too high of a dose may see increased thirst and urination, and edema (swelling, fluid accumulation in the limbs). If you observe any of these signs in your pet, contact your veterinarian immediately.

If your pet experiences an allergic reaction to the medication, signs may include facial swelling, hives, scratching, sudden onset of diarrhea, vomiting, shock, seizures, pale gums, cold limbs, or coma. If you observe any of these signs, contact your veterinarian immediately.

Precautions

Not for use in animals hypersensitive (allergic) to it or drugs similar to it.

Do not use in pregnant or lactating animals (female animals nursing their young).

Do not use in patients with congestive heart failure, kidney disease, or edema (swelling).

Do **NOT** give DOCP intravenously (IV, into the vein).

Consult with your veterinarian regarding the physical examinations and laboratory testing necessary prior to and during treatment with DOCP.

Drug, Food, and Test Interactions

Consult your veterinarian before using vitamins and supplements, diuretics, insulin, digitalis, amphotericin B or potassium-depleting diuretics like Furosemide (Lasix, Salix), since interactions may occur.

Signs of Toxicity/Overdose

May see an increase in drinking and urinating, changes in heart rate or rhythm due to abnormal blood levels of electrolytes, or edema.

Heart enlargement is possible with long-term overdosing.

If you know or suspect your pet has had an overdose, or if you observe any of these signs in your pet, contact your veterinarian immediately.

Keep this and all other medications out of the reach of children and pets.