

DOCTORS FOSTER^{and}SMITH[®] PHARMACY

Patient Information Sheet

PO Box 100, Rhinelander WI 54501
For questions or to consult with a pharmacist, call 1-800-447-3021 Mon-Fri from 9am-6pm CST and Sat from 9am-1pm CST. If you have an emergency situation, please contact your local veterinarian immediately.

Polysulfated Glycosaminoglycan, injectable (PSGAG)

(Adequan)

Common Drug Name

Injectable polysulfated glycosaminoglycan (PSGAG)

dose, skip the one you missed and go back to the regular schedule. Do not give 2 doses at once.

pregnant, or lactating animals (female animals nursing their young).

Common Brand Names

Adequan (injectable)

This medication should only be given to the pet for whom it was prescribed.

Use with caution in animals with kidney or liver disease.

No injectable generic products are available.

Possible Side Effects

With IM injection, few side effects are seen. With intra-articular injection, joint pain, swelling, and lameness may be observed. Rarely, infection of the joint may also occur.

Use with caution in dogs with blood clotting or coagulation disorders.

Storage

Store at room temperature (64-77°F).

Consult with your veterinarian regarding physical examinations and laboratory testing necessary prior to and during treatment with PSGAG.

Uses

Injectable PSGAGs help to control the signs associated with degenerative (non-infectious) or traumatic arthritis in dogs and horses. It has also been used in cats and rabbits.

In dogs, may see a decreased ability for blood to clot, with signs such as bleeding from the nose, blood in the feces, or dark and tarry stools.

Drug, Food and Test Interactions

Consult your veterinarian before using any other medications, including vitamins and supplements, non-steroidal anti-inflammatory drugs (NSAIDs), such as aspirin, etodolac (EtoGesic), deracoxib (Deramaxx), carprofen (Rimadyl), or meloxicam (Metacam), and other drugs affecting blood clotting such as heparin or warfarin, since interactions may occur.

Dose and Administration

Always follow the dosage instructions provided by your veterinarian. If you have difficulty giving the medication, contact your veterinarian.

Consult your veterinarian if you notice any of the above side effects.

This medication is given by intramuscular (IM, into the muscle) injection. Veterinarians may also inject it directly into the joint (intra-articular).

If your pet experiences an allergic reaction to the medication, signs may include facial swelling, hives, scratching, sudden onset of diarrhea, vomiting, shock, seizures, pale gums, cold limbs, or coma. If you observe any of these signs, contact your veterinarian immediately.

Signs of Toxicity/Overdose

Toxicity extremely rare. May see joint pain, swelling, and lameness.

Do not mix this medication with other drugs or chemicals.

Precautions

Not for use in animals hypersensitive (allergic) to this medication.

If you know or suspect your pet has had an overdose, or if you observe any unusual signs, symptoms, or behaviors, contact your veterinarian immediately.

If you miss a dose, give it as soon as you remember. If it is almost time for the next

The manufacturer recommends that this medication not be used in breeding,

Keep this and all other medications out of the reach of children and pets.