

Canine Vaccine Comparison Chart

Drs. Foster & Smith Educational Staff

Vaccinations are an important and easy way to ensure your dog's health. And it's easier than you might think to vaccinate your pet at home. Simply check out our simple to understand [Vaccination Schedule](#) to make an informed decision on what to vaccinate for. Then review our article on [How to Vaccinate Your Dog at Home](#) for a straightforward explanation on the vaccination process.

| Canine Vaccines | Canine Distemper | Hepatitis | Parvovirus | Adenovirus Cough | Parainfluenza | Leptospirosis | Coronavirus | Bordetella | ✓ = Protected against ML = Modified Live K = Killed | | |
|--|------------------|-----------|------------|------------------|---------------|-----------------|-------------|------------|---|---|--------------------|
| COMBINATION VACCINES | | | | | | | | | | | |
| 5-way vaccines | ML | ML | ML | ML | ML | K, 1-4 ** | K | ML | Manufacturer's Recommendations* Min. Age Booster | AVAIL- ABLE IN: | |
| Canine Spectra 5 (Durvet) | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Every 3-4 wks until 16 wks of age (at least 3 doses total) | SINGLE DOSE |
| Duramune Max 5 (Boehringer Ingelheim)+ | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Every 2-3 wks to 12 wks old. Dogs 12 wks or older should receive booster 2-3 wks after initial dose | 25 dose box |
| Nobivac Canine 1-DAPPv (Intervet) ± | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Every 3-4 wks to 12 wks old | 25 dose box |
| Vanguard Plus 5 (Pfizer)+ | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Total of 3 doses, each 3 wks apart | 25 dose box |

| | | | | | | | | | | | |
|--|---|---|---|---|---|------|--|--|-------|--|--------------------|
| Solo-Jec 5 (Boehringer Ingelheim)***+ | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Total of 3 doses, each 3-4 wks apart in puppies 9 wks & younger. Dogs 12 wks or older should receive booster 2-3 wks after initial dose. | SINGLE DOSE |
| 5-way vaccines plus coronavirus | | | | | | | | | | | |
| Duramune Max 5-CvK (Boehringer Ingelheim)+ | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Every 2-3 wks to 12 wks old. Dogs 12 wks or older should receive booster 2-3 wks after initial dose | 25 dose box |
| Vanguard Plus 5/CV (Pfizer)+ | ✓ | ✓ | ✓ | ✓ | ✓ | | | | 6 wks | Total of 3 doses, each 3 wks apart | 25 dose box |
| 9-way vaccines | | | | | | | | | | | |
| Duramune Max 5/4L (Boehringer Ingelheim)+ | ✓ | ✓ | ✓ | ✓ | ✓ | 4 ** | | | 6 wks | Every 2-3 wks to 12 wks old | 25 dose box |
| Univac 9 (Agrilabs)***+ | ✓ | ✓ | ✓ | ✓ | ✓ | 4 ** | | | 6 wks | Total of at least 3 doses, each 3-4 wks apart until 16 wks of age. | SINGLE DOSE |
| Vanguard Plus 5 L4 (Pfizer)+ | ✓ | ✓ | ✓ | ✓ | ✓ | 4 ** | | | 6 wks | Total of 3 doses, each 3 wks apart | 25 dose box |

| | | | | | | | | | | | |
|--|---|---|---|---|---|-----|---|---|-------|--|--------------------|
| Canine Spectra 9 (Durvet)***+ | ✓ | ✓ | ✓ | ✓ | ✓ | 4** | | | 6 wks | Every 2-4 wks to 16 wks old. | SINGLE DOSE |
| 9-way vaccines plus coronavirus | | | | | | | | | | | |
| Duramune Max 5-CvK/4L (Boehringer Ingelheim)+ | ✓ | ✓ | ✓ | ✓ | ✓ | 4** | ✓ | | 6 wks | Every 2-3 wks to 12 wks old | 25 dose box |
| Parvovirus and/or coronavirus vaccines | | | | | | | | | | | |
| Duramune Max PV (Boehringer Ingelheim)+ | | | ✓ | | | | | | 6 wks | Every 2-3 wks to 12 wks old. Dogs over 12 wks should initially receive 1 dose and a 2nd dose 2-3 wks later | 25 dose box |
| Duramune Cv-K (Boehringer Ingelheim) | | | | | | | ✓ | | 6 wks | Every 2-3 wks to 12 wks old | 25 dose box |
| Vanguard Plus CPV (Pfizer)+ | | | ✓ | | | | | | 6 wks | Total of 3 doses, each 3 wks apart | 10 dose vial |
| Nobivac Puppy-DPv (Intervet)+ | ✓ | | ✓ | | | | | | 4 wks | Every 3-4 wks to 12 wks old | 25 dose box |
| Bordetella and/or parainfluenza | | | | | | | | | | | |
| Bronchicine CAe (Pfizer)+ | | | | | | | | ✓ | 8 wks | Total of 2 doses, each 2-4 wks apart | 10 dose vial |
| Nobivac Intra-Trac KC (Intervet) NOSE DROPS - NOT INJECTABLE | | | | | ✓ | | | ✓ | 3 wks | Every 12 mths | 10 dose vial |

| | | | | | | | | | | | |
|---|----|--|--|---|---|--|--|---|-------|---|--------------------|
| Nobivac Intra-Trac3 ADT (Schering)+ NOSE DROPS - NOT INJECTABLE | | | | ✓ | ✓ | | | ✓ | 3 wks | Every 12 mths | 25 dose box |
| Kennel Jec-2 (Durvet)+ NOSE DROPS - NOT INJECTABLE | | | | | ✓ | | | ✓ | 3 wks | Every 12 mths. Puppies vaccinated between 3-6 wks should be re-vaccinated at 6 wks. | SINGLE DOSE |
| SINGLE PROTECTION VACCINES | | | | | | | | | | | |
| Lyme vaccine | | | | | | | | | | | |
| Lymevox (Pfizer) | NA | | | | | | | | 9 wks | 2-3 wks after initial dose | 10 dose vial |

*These are the manufacturer's recommendations for the age at when to start vaccinating and the booster schedule. Your veterinarian may have different recommendations for minimum age, boosters, and annual vaccinations based on the breed, health status, and potential of the dog to be exposed to the disease. Many veterinarians do not recommend the use of a leptospirosis vaccine (7-way or 9 -way) in dogs less than 12 weeks of age. See [Vaccination Schedule for Dogs](#).

**Number of strains included.

***Residents of CT, DE, IL, NJ, or NY, please ask your veterinarian for a prescription. Residents of NY may use a certificate of need. Needle and syringe are included with vaccine.

+Residents of MN, please ask your veterinarian for a prescription.